



<b>Mercury Toxicity and Lichen Planus</b>	<b>Clinical Protocol</b>
<b>Oral Lichen Planus and other oral lesions including squamous cell cancer: The Primary Cause is Immune Reactivity to Amalgam Fillings</b>	
<ul style="list-style-type: none"><li>• Mercury is one of the most toxic substances in existence and is known to bioaccumulate in the body of people and animals that have chronic exposure (35).</li><li>• Mercury from occupational exposure and dental fillings is primarily from elemental mercury vapor.</li><li>• Mercury vapor is highly absorbed by the lungs and in saliva or blood is rapidly converted to ionic or methyl mercury.</li><li>• Mouth bacteria and yeast as well as other methyl donors convert other forms of mercury to <a href="#">methyl mercury</a>, so that most mercury in the blood is methyl mercury regardless of source (9, 42).</li><li>• Mercury in amalgam fillings, because of its high volatility and galvanic action due to presence of dissimilar metals in the mouth, has been found to be continuously vaporized and also released into the body through galvanic currents (29, 43, etc.), and has been found to be the largest source of mercury in the majority of people WHO (27), 9, 30, 32, 42, 1, 14).</li><li>• The level of daily exposure commonly exceeds the U.S. EPA health guideline for daily mercury exposure (35, 42).</li><li>• Mercury vapor given off by amalgam fillings accumulates in the teeth, tooth roots, gums, jawbone, and oral tissue. The number of amalgam surfaces has a statistically significant correlation to the level of mercury in oral mucosa and saliva (1, 12, 13, 26, 30, 33, 36, and 42).</li><li>• High levels of mercury have been documented to accumulate in the gums, jawbone, and oral mucosa of those with amalgam fillings and to be transferred to the blood stream and other parts of the body (43).</li><li>• Concentrations of mercury in oral mucosa for a population of patients with 6 or more amalgam fillings taken during oral surgery were 20 times the level of controls (25).</li><li>• Studies have shown mercury travels from amalgam into dentin, root tips, and the gums, with levels in roots tips as high as 41 parts per million (ppm) (25).</li><li>• Studies have shown that mercury in the gums, such as from root caps for root canal teeth or amalgam tattoos, result in chronic inflammation and proliferation of inflammatory cells, in addition to migration to other parts of the body (31, 7, 6, 43, and 151).</li><li>• Mercury, silver, and other metals from fillings can be seen in the tissues as amalgam “tattoos”, which have been found to accumulate in the oral mucosa as granules along collagen bundles, blood vessels, nerve sheaths, elastic fibers, membranes, striated muscle fibers, and acini of minor salivary glands. Dark granules are also present intracellularly within macrophages, multinucleated giant cells, endothelial cells, and fibroblasts, and metals also accumulate in tooth roots and the jaw bone (7, 6).</li></ul>	

- There is, in most cases, chronic inflammatory response or macrophagic reaction to the metals (7, 18), usually in the form of a foreign body granuloma with multinucleated giant cells of the foreign body and Langhans types (29).
- In a group of patients with amalgam tattoos that were tested, 74% of the patients revealed high lymphocyte reactivity (positive MELISA test) to one or more metal components of dental restorations(7k). The majority of MELISA positive patients suffered from serious health problems (various allergies, autoimmune diseases, Parkinson's syndrome etc.).
- Nickel and inorganic mercury were the most common sensitizers in vitro. The cytokine assay revealed that mercury chloride activated predominantly TH2 lymphocytes, while nickel chloride activated mainly TH1 lymphocytes.
- Many dentists are not aware that the main source of amalgam tattoos is “oral galvanism”, where electric currents caused by mixed metals in the mouth take the metals into the gums and oral mucosa, accumulating at the base of teeth with large fillings or metal crowns over amalgam base (29, 43). Such mercury, including that in the commonly formed amalgam tattoos, moves to other parts of the body over time in significant amounts and more rapidly than the other metals.
- Macrophages remove mercury by phagocytosis and the mercury moves to other parts of the body through the blood and along nerves (7).
- Another study (7l) demonstrated a dense mononuclear inflammatory infiltrate associated with large and powdered debris and positivity for HLA-DR and MT in inflammatory cells. While blood vessel walls and connective fibers impregnated with powdered particles were negative for HLA-DR, they were positive for MT. In addition, wherever epithelial basement membrane impregnation by powdered amalgam particles was observed, a strong positivity for MT was detected.
- These findings demonstrate that residual elements of AT still have noxious local effects over tissues. Such metals are documented to commonly cause local and systemic lesions along with other health effects, which usually recover after removal of the amalgam tattoo by surgery (7fghim). The high levels of accumulated mercury also are dispersed to other parts of the body (43).
- The amount of mercury in saliva averaged between 1.5 to 1.9 micrograms per liter for each amalgam filling (30ab), enough to cause daily exposure of 10 to 100 micrograms of mercury.
- The amount of mercury released by a gold alloy bridge over amalgam over a 10 year period was measured to be approx. 101 milligrams (mg) (60% of total) or 30 micrograms (ug) per day (1), and other studies have found similar results (26, 42).
- The average mercury levels in gum tissue near amalgam fillings are often over 100 ppm (29), and levels in oral mucosa removed during oral surgery averaged over 2 ppm (over 20 times controls) and levels in root tips of 41 ppm (25, 29, 7).
- Having dissimilar metals in the teeth (e.g. gold and mercury) causes galvanic action, electrical currents, and much higher mercury vapor levels and mercury levels in tissues. (26, 28, 29, 1, 2, 4, 5, 7, 8, 25).
- The level of mercury in the gums or jaw bone is often 1000 ppm near a gold cap on an amalgam filling (5, 3, 6, 8, 10), and similar levels as high as 5600 ppm have been found in the jaw bone under large amalgam fillings or gold crowns over amalgam by German oral surgeons (44).

- These levels are among the highest levels ever measured in tissues of living organisms, exceeding the highest levels found in chronically exposed chloral kali workers, those who died from mercury in Minamata, or animals that died from mercury poisoning.
- The FDA action level for warnings of dangerous levels in fish or food is 1 ppm and the EPA health criterion level is 0.3 ppm.
- Amalgam also releases significant amounts of silver, tin, and copper which also have toxic effects, with organic tin compounds formed in the body being even more neurotoxic than inorganic mercury.
- Toxic/allergic reactions to toxic metals such as mercury often result in autoimmune conditions, such as lichen planus lesions in oral mucosa or gums, and play a roll in pathogenesis of periodontal disease.
- Oral lichen planus has been found to be an autoimmune process in which the Immune Th1 T-cells mediate the reactivity, including Lymphotoxin-alpha (LTa), Tumor Necrosis Factor-alpha (TNFa), and Interferon-gamma (IFNa) (18, 37, 40b).
- A high percentage of patients with oral mucosal problems (37, 18), along with other autoimmune conditions such as chronic fatigue (23, 39), MS or lupus (40), have significant immune reactions to mercury, palladium, gold, and nickel (37, 23).
- **Removal of amalgam fillings usually led to cure or significant improvement for oral lichen planus** (15-17, 20-22, 24, 37, etc.), as well as **for oral keratosis** (pre-cancer) (16b, 45), and **most of other oral health problems** including metallic taste, tender teeth, mouth sores, bad breath, bleeding gums and throat irritation (43). A connection between mercury immune reactivity from amalgam and oral cancers has also been demonstrated (18, 19).
- Most cases of CFS, MS, or lupus patients also had significant immune reactions to inorganic mercury (MELISA test) and removal of amalgam fillings usually results in cure or significant improvement of such conditions (23, 39, 40, and 11).
- In one clinic (21) that replaced amalgams for a large number of such patients, there was cure or significant improvement in over 90% of cases.
- A Jerome meter was used to measure mercury vapor level in the mouth, and many had over 50 micrograms mercury per cubic meter of air, far above the Government health guideline for mercury (35).
- In a recent study of patients with OLP, 60% showed sensitization to 1 or more allergens using a patch test (17a). The greatest frequency of positive reactions was to dental metals.
- The order of tested metals, according to frequency of positive reactions, was mercury, amalgam nickel, palladium, cobalt, gold, chrome, and indium. However, patch tests have been found to not be a reliable indicator of mercury immune reactivity or allergy.
- In large number of clinical trials by doctors treating OLP, between 39 and 53% of patients tested by patch tests were indicated to be reactive to mercury (16abc, 17, and 24a). However, when patients had amalgams replaced, the majority recovered or significantly improved in a relatively short time period regardless of patch test results (15, 16abc, 17, 24, and 37). Thus, the authors recommend replacement of amalgam in all cases of OLP and similar conditions.
- The MELISA blood lymphocyte immune reactivity test appears to be a more accurate indicator of immune reactivity than the patch test (37, 39, and 40). When patch tests are to be used it should be noted that the clinical trials found that mercury immune reactivity

- is often a delayed reaction, with positive patch test observed only later on the 10th or 17<sup>th</sup> day of the test (17, and 24a).
- Patients with OLP also commonly have been found to be immune reactive to gold or nickel (17, 16d, 37, and 40) so that replacement of gold or nickel crowns may be beneficial in such patients when amalgam replacement is not sufficient to resolve the problem.
- Oral lichen planus and oral lesions, caused most commonly by reactivity to mercury, are inflammatory pre-cancerous conditions that have been well documented in the literature to often develop into oral squamous cell carcinoma (OSCC) (46, and 90a).
- Infection and chronic inflammation have been found to contribute to carcinogenesis through inflammation-related mechanisms (47 and 48). Inflammatory bowel diseases are associated with colon carcinogenesis and inflammatory oral conditions such as oral lichen planus (OLP) and leukoplakia are associated with OSCC.
- Previous studies have shown significant increases of NF-kappaB dependent cytokines, Tumor Necrosis Factor-alpha (TNF-a), IL-1alpha, IL-6, and IL-8 in different oral fluids from oral lichen planus (OLP) patients (48).
- In samples of whole, unstimulated saliva, in this study, for moderate and severe OLP dysplasia, the level of each cytokine was significantly higher than in control.
- In moderate dysplasia, TNF-alpha and IL-1alpha were significantly increased at a level without difference from OSCC, but IL-6 and IL-8 was detected at a concentration significantly lower than OSCC.
- In severe OLP dysplasia, the level of TNF-alpha was not significantly different from that of Oral Squamous Cell Carcinoma.
- The study confirmed preclinical data that NF-kappaB dependent cytokines are up-regulated in pre-malignant OLP and oral carcinogenesis.
- Cytologic and DNA-cytometric examination of oral lesions and oral lichen planus have also been found to be reliable indicators of OLP cases becoming malignant (49). Immunolabeled oncoproteins were found to be modified in the premalignant leukoplakia, oral lichen planus and in squamous cell cancer (49, and 46a).
- The evidence supports that dental amalgam is the most common cause of oral squamous cell carcinoma, similar to the fact it is the most common cause of OLP. The available pretreatment dental records of 133 patients with carcinoma of the tongue seen at the British Columbia Cancer Agency between 1958 and 1992 were reviewed. The majority had amalgam fillings on the side of the tongue involved in the carcinoma. Of the 7 patients with amalgams on only one side of the mouth, 6 cases of oral cancer had amalgams on the side of the cancer and only 1 on the side without amalgams (50).
- People with oral lichen planus often develop OLP at multiple sites (51) and also can have lichen planus in other locations such as the esophagus (52) or genitals (53).
- In one study, 41 women diagnosed with OLP underwent gynecological exam and 75.6% were found to have evidence of genital involvement, vulvar lichen planus or vulvar lichen sclerosis (53b). Such inflammatory conditions can also become cancerous (52, and 54).
- Two siblings with long standing cutaneous lichen planus of the esophagus both developed squamous cell carcinoma (52). Since immune reactivity to mercury is the most common cause of OLP and OSCC and since immune reactivity to mercury is a systemic condition (37, etc.), systemic immune reactivity to mercury might be the most likely cause of lichen planus and resulting squamous cell cancers of other organs such as the esophagus and genitals.

## References

- Windham, B. (Ed), **Oral Lichen Planus and other oral lesions including squamous cell cancer: The Primary Cause is Immune Reactivity to Amalgam Fillings.**
- Vimy, M.J., Lorscheider, F.L. **Intra oral Mercury released from dental amalgams and estimation of daily dose;** J. Dent Res. 64(8):1069-1075, 1985.
- B Momoi Y, et al; **Measurement of galvanic current and electrical potential in extracted human teeth,** J Dent Res, 65(12): 1441-1444;  
Holland, R.I., **Galvanic currents between gold and amalgam.** Scand J Dent Res, 1980, 88:269-72;  
Wang Chen C.P. and Greener, E.H., **A galvanic study of different amalgams,** Journal of Oral Rehabilitation, 1977, 4:23-7;  
Lemons, J.E. et al, **Intraoral corrosion resulting from coupling dental implants and restorative metallic systems,** Implant Dent, 1992, 1(2):107-112.
- C. Malmstrom et al., **"Silver amalgam: an unstable material"**, Swedish paper translated in Bio-Probe Newsletter, Vol 9(1):5-6, Jan. 1993, <http://home.swipnet.se/misac/research1.html> & Tidsskrift for Tandlaeger, Oct 1989, 5:85-87, <http://home.swipnet.se/misac/research1.html>.
- Raue, H., **"Resistance to therapy; Think of tooth fillings"**, Medical Practice, vol. 32, n.72, p.2303- 2309, 6 Sept 1980.
- Till et al. Zahnarztl. Welt/reform 1978:87;1130-1134  
Olsson, S. et al, **"Release of elements due to electrochemical corrosion of dental amalgam"** J of Dental Research, 1994, 73:33-43;  
Fusayama, T. et al, J Dental Res, 1963, 42:1183-1197;  
Tuija, P. , Yli-Urpo, A., **A histological study of the influence of galvanic current between metal alloys in connective tissue in the rat,** 1973, Proc Fin J Den Soc 69:1-6;  
Horsted-Binslev, P., Danscher, G. **Dentinal and pulpal uptake of mercury from lined and unlined amalgam restorations.** Eur J Oral Sci 1997, 105: 338-43.
- Huggins, H.A., Levy, T.E., **Uniformed Consent: the hidden dangers in dental care,** 1999, Hampton Roads Publishing Company Inc; & Huggins, H.A., Its All in Your Head, 1997 & Center for Progressive Medicine, 1999, <http://www.hugnet.com>.
- Buchner, A. et al, **"Amalgam tattoo of the oral mucosa: a clinicopathologic study of 268 cases"**, Surg Oral Med Oral Pathol, 1980, 49(2):139-47;  
Owens, B.M., Johnson, W.W., Schuman, N.J. **Oral amalgam pigmentations (tattoos): a retrospective study.** Quintessence Int. 1992 Dec; 23(12):805-10.  
Forsell, M. et al, **Mercury content in amalgam tattoos of human oral mucosa and its relation to local tissue reactions.** Euro J Oral Sci 1998; 106(1):582-7;  
Harrison, J.D. et al, **Amalgam tattoos: light and microscopy and electron-probe micro-analysis;**  
Kanzaki, T. et al, **Electron microscopic X-ray microanalysis of metals deposited in oral mucosa.** J Dermatol 1992; 19(8):487-92;

Nilner, K. et al, **In vitro testing of dental materials by means of macrophage cultures.** J Biomed Mater Res 1986;20(8):1125-38;

Weaver, T., Auclair, P.L.; **Amalgam tattoo as a cause of local and systemic disease?** Oral Surg. Oral Med. Oral Pathol. 1987;63:137-40;

Rusch-Behrend, G.D., Gutmann, J.L. **Management of diffuse tissue argyria subsequent to endodontic therapy: report of a case.** Quintessence Int. 1995 Aug;26(8):553-7;

Mayall, F.G., Hickman, J., Knight, L.C., Singharo, S. **An amalgam tattoo of the soft palate: a case report with energy dispersive X-ray analysis.** J Laryngol Otol. 1992 Sep; 106(9):834-5;

Kissel, S.O., Hanratty, J.J. **Periodontal treatment of an amalgam tattoo.** Compend Contin Educ Dent. 2002 Oct;23(10):930-2, 934, 936;

**Amalgam-tattoo-associated oral lichenoid lesion.** Staines, K.S., Wray, D. Contact Dermatitis. 2007 Apr;56(4):240-1;

**In vivo effects of dental casting alloys.** Neuro Endocrinol Lett. 2006 Dec;27 Suppl 1:61-8. Venclíková, Z., Benada, O., Bártová, J., Joska, L., Mrklas, L., Procházková, J., Stejskal, V., Podzimek, S.;

**Immunolocalization of HLA-DR and metallothionein on amalgam tattoos.** Braz Dent J. 2004;15(2):99-103. Epub 2005 Mar 11, Leite, C.M., Botelho, A.S., Oliveira, J.R., Cardoso, S.V., Loyola, A.M., Gomez, R.S., Vaz, R.R.;

**Chronic inflammation and pain inside the mandibular jaw and a 10-year forgotten amalgam filling in an alveolar cavity of an extracted molar tooth.** Ultrastruct Pathol. 2005 Sep-Oct;29(5):405-13, Kaufmann, T., Bloch, C., Schmidt, W., Jonas, L.

- Muller, A.W., Van Loon, L.A., Davidson, C.L. **Electrical potentials of restorations in subjects without oral complaints.** J Oral Rehabil. 1990 Sep;17(5):419-24;

Inovay, J., Banoczy, J. (1961)**The role of electrical potential differences in the etiology of chronic diseases of the oral mucosa.** Journal of Dental Research, 40, 884;

Arvidson,, K., **Corrosion studies of dental gold alloy in contact with amalgam,** Swed. Dent. J 68: 135-139,1984;

Lemons, J.E. et al, **Interoral corosion resulting from coupling dental implants and restorative metallic systems,** Implant Dent, 1992, 1(2):107-112;

Skinner, E.W., **The Science of Dental Materials,** 4th Ed.revised, W.B.Saunders Co., Philadelphia, p 284-285,1957;

- Kingman, A., Albertini, T., Brown, L.J. National Institute of Dental Research, **Mercury concentrations in urine and blood associated with amalgam exposure in the U.S. military population,** J Dent Res. 1998 Mar;77(3):461-71.
- Freden, H. et al, **Mercury in gingival tissues adjacent to amalgam fillings,** Odontol Revy, 1974, 25(2): 207-210; H Reden, Odontol Revy, 25,1971,207-210.
- Katsunuma et al, **Anaphylaxis improvement after removal of amalgam fillings,** Annals of Allergy, 1990, 64(5):472-75;

Drouet, M. et al, **Is mercury a respiratory tract allergen?** Allerg Immunol (Paris),1990,22(3):81.

- Skare, I., **Mass Balance and Systemic Uptake of Mercury Released from Dental Fillings, Water, Air, and Soil Pollution**, 80(1-4):59-67, 1995;
- Skare, J., Enqvist, A., **Human exposure to mercury and silver released from dental amalgam restorations**. Archives of Env Health 1994; 49(5): 384-94.
- Bjorkman, L. et al, **Mercury in Saliva and Feces after Removal of Amalgam Fillings**, J Dent Res 75: 38- IADR Abstract 165, 1996; Toxicology and Applied Pharmacology, May 1997, 144(1), p156-62.
- Begerow, J. et al, **Long-term mercury excretion in urine after removal of amalgam fillings**, Int Arch Occup Health 66:209-212, 1994.
- **Association between oral lichenoid reactions and amalgam restorations**, Pezelj-Ribarić, S., Prpić, J., Miletić, I., Brumini, G., Soskić, M.S., Anić, I.; J Eur Acad Dermatol Venereol. 2008 Nov;22(10):1163-7. Epub 2008 Apr 3;
- Dunsche, A. et al, **Oral lichenoid reactions associated with amalgam: improvement after amalgam removal**. British Journal of Dermatology 2003 Jan;148:1:70-6;  
Smart, E.R. et al, **Resolution of lichen planus following removal of amalgam restorations**, Br Dent J 178(3): 108-112,1995;  
Markow, H., **Regression from urticaria following dental filling removal**.,New York State J Med,1943: 1648-1652;  
Sasaki, G. et al, **Three cases of oral lichenosis caused by metallic fillings**, J. Dermatol, 23 Dec, 1996; 12:890-892.
- Skoglund. A., Scand J Dent Res 102(4): 216-222, 1994; and 99(4):320-9,1991;  
Ostman, P.O. et al, **Clinical & histologic changes after removal of amalgam** Oral Surgery, Oral Medicine, and Endodontics, 1996, 81(4):459-465;  
Wong, L. and Freeman, S., **Oral lichenoid lesions (OLL) and mercury in amalgam fillings**, Contact Dermatitis, Vol 48 Issue 2 Page 74 - February 2003;  
Alanko, K., Kanerva, L., Jolanki, R., Kannas, L., Estlander, T., **Oral mucosal diseases investigated by patch testing with a dental screening series**. Contact Dermatitis. 1996 Apr;34(4):263-7.
- **Oral lichenoid lesions and allergy to dental materials**. Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub. 2007 Dec;151(2):333-9. Ditrichova,, D., Kapralova, S., Tichy, M., Ticha, V., Dobesova, J., Justova, E., Eber, M., Pirek, P.  
Koch, P. et al, **Oral lichenoid lesions,mercury hypersensitivity, ...**, Contact Dermatitis, 1995, 33(5):323-328.  
Freeman, S. et al, **Oral lichenoid lesions caused by allergy to mercury in amalgam**, Contact Dermatitis,1995,33(6): 423-7(Denmark).  
Mobacken, H. et al, Contact Dermatitis,1984,10:11.
- **Expression of lymphotoxin-alpha by keratinocytes: a further mediator for the lichenoid reaction**. Middel, P., Lippert, U., Hummel, K.M., et al, Pathobiology. 2000;68(6):291-300.
- Hougeir, F.G., Yiannias, J.A., Hinni, M.L., Hentz, J.G., el-Azhary, R.A., **Oral metal contact allergy: a pilot study on the cause of oral squamous cell carcinoma**. Int J Dermatol. 2006 Mar;45(3):265-71.

- Berglund, F., **Case reports spanning 150 years on the adverse effects of dental amalgam**, Bio-Probe, Inc., Orlando, FL, 1995; ISBN 0-9410011-14-3 (245 cured)
- Lichtenberg, H., **Symptoms before and after proper amalgam removal in relation to serum-globulin reaction to metals**, Journal of Orthomolecular Medicine, 1996, 11(4): 195-203. (119 cases)
- Henriksson, E. et al, **Healing of Lichenoid Reactions following Removal of Amalgam**, J Clinical Periodontol, V22, N4, p287-94, 1995.  
Forsbec, M. et al, Journal of Clinical Immunology, 16(1):31-40, Jan 1996.
- Tibbling, L. et al, **Immunological and brain MRI changes in patients with suspected metal intoxication**, Int J Occup Med Toxicol 4(2):285-294, 1995.
- Laine, J. et al, **Resolution of oral lichenoid lesions after replacement of amalgam restorations**, Br J Dermatol, 1992, 126(1):10-15;  
Ibbotson, S.H. et al, **The relevance of amalgam replacement on oral lichenoid reactions**, British Journal of Dermatology, 134(3):420-3, 1996.
- Willershausen, B. et al, **Mercury in the mouth mucosa of patients with amalgam fillings**, Dtsch Med Wochenschr, 1992, 117:46, 1743-7.
- Pleva, J., Orthomol. J., Psych, Vol 12, No.3, 1983 & J. Of Orthomol. Medicine 1989, 4:141- 148. & **Mercury- A Public Health Hazard**, Reviews on Environmental Health, 1994, 10:1-27.
- World Health Organization (WHO), 1991, Environmental Health criteria 118, Inorganic Mercury, WHO, Geneva. Envir. H. Crit. 101, Methyl Mercury.  
Halbach, 1995, **estimation of mercury dose ...**, Int. Archives of Occupational & Environmental Health, 67: 295-300.  
Sandborgh-Englund, G., **Pharmakinetics of mercury from dental amalgam**, Gotab (Stolckholm), 1998, 1-49.
- Brune, D. et al, Scand J Dent Res, 1983, 19:66-71 & Sci Tot Envir, 1985, 44:....; **Metal release from dental materials**, Biomaterials, 1986, 7, 163-175.
- Nogi, N., **Electric current around dental metals as a factor producing allergic metal ions in the oral cavity**, Nippon Hifuka Gakkai Zasshi, 1989, 99(12):1243-54;  
Rose, M.D. et al, Eastman Dental Institute, **The tarnished history of a posterior restoration**, Br Dent J 1998; 185(9):436;  
Certosimo, A.J. et al, National Naval Dental Center, **Oral Electricity**, Gen Dent, 1996, 44(4):324-6.  
Ogletree, R.H. et al, School of Materials Science, GIT, Atlanta, **Effect of mercury on corrosion of eta Cu-Sn phase in dental amalgams**, Dent Mater, 1995, 11(5):332-6.  
Meyer, R.D. et al, **Intraoral galvanic corrosion**, Prosthet Dent, 1993, 69(2):141-3.  
Owens, B.M. et al, **Localized galvanic shock after insertion of an amalgam restoration**, Compenium, 1993, 14(10), 1302, 1304, 1306-7.  
Johansson, E., Liliefors, T., **Heavy elements in root tips from teeth with amalgam fillings**, Department of Radiation Sciences, Division of Physical Biology, Box 535, 751 21 Uppsala, Sweden.  
Cheshire, William P., Jr. **The shocking tooth about trigeminal neuralgia**. New England Journal of Medicine, Vol. 342, June 29, 2000, p. 2003.

- Kraub, P. Dr. & Deyhle, M., Universitat Tubingen- Institut fur Organische Chemie, **Field Study on the Mercury Content of Saliva**, 1997  
<http://www.uni-tuebingen.de/KRAUSS/amalgam.html>.  
Monaci, F., Bargagli, E., Bravi, F., Rottoli, P., **Concentrations of major elements and mercury in unstimulated human saliva**. Biol Trace Elem Res. 2002 Dec;89(3):193-203.
- Nadarajah, V. et al, **Localized cellular inflammatory response to subcutaneously implanted dental mercury**, J Toxicol Environ Health, 1996, 49(2):113-25;  
Kulacz & Levy, "The Roots of Disease". Xlibris Corporation at 1-888-795-4274  
[www.xlibris.com](http://www.xlibris.com).
- Richardson, Mark, Environmental Health Directorate, Health Canada. **Assessment of Mercury Exposure and Risks from Dental Amalgam**, 1995, Final Report.
- Vimy, M.J. and Lorscheider, F.L., Faculty of Medicine, Univ. Of Calgary, J. Trace Elem. Exper. Med., 1990, 3: 111-123.
- Ziff, M.F., **Documented clinical side effects to dental amalgams**, ADV. Dent. Res., 1992; 1(6):131-134.  
Ziff, S., **Dentistry without Mercury**, 8th Edition, 1996, Bio-Probe, Inc., ISBN 0-941011-04-6; **Dental Mercury Detox**, Bio-Probe, Inc. (Cases: FDA Patient Adverse Reaction Reports-762, Hanson, M. Dr.-Swedish patients-519, Lichtenberg, H. Dr.-100 Danish patients, Larosek, P. Dr.- 80 Canadian patients, Sibley, R. Dr.-86 Colorado patients, Dr. A.V.Zamm, 22 patients) <http://www.flcv.com/hgrecovp.html>.
- Agency for Toxic Substances and Disease Registry, U.S. Public Health Service, **Toxicological Profile for Mercury**, 1999; & Apr 19,1999 Media Advisory, New MRLs for toxic substances, MRL:elemental mercury vapor/inhalation/chronic. MRL: methy mercury/ oral/acute; & <http://www.atsdr.cdc.gov/mrls.html>.
- Dauserer, M., **Improvement of Nerve and Immunological Damages after Amalgam Removal**, Amer. J. Of Probiotic Dentistry and Medicine, Jan 1991.
- Stejskal, V.D., Forsbeck, M., Cederbrant, K.E., Asteman, O.. **Mercury-specific lymphocytes: an indication of mercury allergy in man**. J Clin Immunol. 1996 Jan; Vol 16(1):31-40; [www.melisa.org](http://www.melisa.org).
- Tosti, A. et al, **Contact stomatitis**, Semin Cutan Med Surg, 1997, 16(4):314-9.  
Nakada, T. et al, **Patch test materials for mercury allergic contact dermatitis**, Dermatitis, 1997, 36(5):237-9.  
Guttman-Yassky, E., Weltfriend, S., Bergman, R., **Resolution of orofacial granulomatosis with amalgam removal**. J Eur Acad Dermatol Venereol. 2003 May;17(3):344-7.  
Lazarov, A., Kidron, D., Tulchinsky, Z., Minkow, B., **Contact orofacial granulomatosis caused by delayed hypersensitivity to gold and mercury**. J Am Acad Dermatol. 2003 Dec;49(6):1117-20.
- Sterzl, I., Prochazkova, J., Stejskal, V.D.M. et al, **Mercury and nickel allergy: risk factors in fatigue and autoimmunity**. Neuroendocrinology Letters 1999; 20:221-228.  
Stejskal, V., Hudecek, R., Mayer, W., **Metal-specific lymphocytes: risk factors in CFS and other related diseases**, Neuroendocrinology Letters, 1998 ; [www.melisa.org](http://www.melisa.org).

- Prochazkova, J., Sterzl, I., Kucerova, H., Bartova, J., Stejskal, V.D.; **The beneficial effect of amalgam replacement on health in patients with autoimmunity.** Neuro Endocrinol Lett. 2004 Jun;25(3):211-8; <http://www.melisa.org/pdf/Mercury-and-autoimmunity.pdf>.

Stejskal, J., Stejskal, V., **The role of metals in autoimmune diseases and the link to neuroendocrinology** Neuroendocrinology Letters, 20:345-358, 1999

- Windham, B.(Ed)., Anotated Bibliography: **Exposure Levels and Health Effects of Mercury from Amalgam Fillings and Results of Filling Replacement**, 2000,(over 4000 Medical Studies & scientific journal references and 60,000 clinical cases of replacement followed by doctors); [www.flcv.com/amalg6.html](http://www.flcv.com/amalg6.html).
- Windham, B.(Ed), Annotated bibliography: **Level of Exposure to inorganic and methyl mercury from dental amalgam**, 2005; [www.flcv.com/damspr1.html](http://www.flcv.com/damspr1.html).
- Windham, B.(Ed), **Accumulation of mercury in the oral cavity from dental amalgam and oral effects**; [www.flcv.com/periodon.html](http://www.flcv.com/periodon.html).
- Schiwara, H.W., (Medical Laboratory) Arzte fur Laboratoriumsmedizen, D-28357 Bremen; & Heavy Metal Bul, 1999, 1:28.
- Omura, Y. et al, Heart Disease Research Foundation, NY,NY, **Role of mercury in resistant infections and recovery after Hg detox with cilantro**, Acupuncture & Electro-Therapeutics Research, 20(3):195-229, 1995. **Mercury exposure from silver fillings**, Acupuncture & Electrotherapy Res, 1996, 133
- **Evaluation of proliferative potential in oral lichen planus and oral lichenoid lesions using immunohistochemical expression of p53 and Ki67**, Acay, R.R., Felizzola, C.R., et al, Oral Oncol. 2006 May; 42 (5):475-80. Epub 2005 Dec 20.

**Premalignant nature of oral lichen planus**, Laeijendecker, R., van Joost, T. et al, Acta Derm Venereol. 2005;85 (6):516-20.

**Oral cancer development in patients with oral lichen planus**, Barnard, N.A., Scully, C., et al, J Oral Pathol Med. 1993 Oct;22(9):421-4.

**The possible premalignant character of oral lichen planus and oral lichenoid lesions: A prospective five-year follow-up study of 192 patients**, van der Meij, E.H., Mast, H., van der Waal, I., Oral Oncol. 2006 Nov 15.

**Squamous cell carcinoma of the tongue arising in lichen planus: a case report and review of the literature**, Camisa, C., Hamaty, F.G., Gay, J.D., Cutis. 1998 Oct;62 (4):175-8.

**Oral squamous cell carcinoma arising in a patient with long-standing lichen planus.** A case report, Katz, R.W., Brahim, J.S., Travis, W.D. Oral Surg Oral Med Oral Pathol. 1990 Sep;70(3):282-5.

**Squamous cell carcinoma on the dorsum of the tongue arising in a long-standing lesion of erosive lichen planus**, Fowler, C.B., Rees, T.D., Smith, B.R., J Am Dent Assoc. 1987 Nov;115(5):707-10.

**Daily practice: appreciation of the oncologic risk of oral lichen planus**, Castermans, Elias S., Castermans, A., Rev Belge Med Dent. 1975;30(1):33-8.

**Epithelioma on oral lichen planus (apropos of 40 new cases)**; Cernea, P., Kuffer, R., Brocheriou, C., Actual Odontostomatol (Paris). 1971 Dec;96:473-90.

**Cancer development in oral lichen planus.** A follow-up study of 327 patients, Fulling, H.J.. Arch Dermatol. 1973 Nov;108 (5):667-9.

**Carcinoma arising in erosive lichen planus in the midline of the dorsum of the tongue**, Pogrel, M.A., Weldon, L.L., Oral Surg Oral Med Oral Pathol. 1983 Jan;55(1):62-6.

**The association of chronic inflammatory disease in lichen planus with cancer of the oral cavity**, Deeb, Z.E., Fox, L.A., deFries, H.O.. Am J Otolaryngol. 1989 Sep-Oct;10 (5):314-6.

**Squamous cell carcinoma of the oral cavity-chronic oral ulcerative disease as a possible etiologic factor**, Faraci, R.P., Schour, L., Graykowski, E.A., J Surg Oncol. 1975;7(1):21-6.

**Oral lichen planus and squamous carcinoma**. Case report and update of the literature, Kaplan, B., Barnes, L., Arch Otolaryngol. 1985 Aug;111(8):543-7.

**Oral lichen planus and squamous carcinoma: case report and update of the literature**, Kaplan, B.R., R I Dent J. 1991 Winter;24(4):5-9, 11-4;

- **Oxidative and nitrative DNA damage in animals and patients with inflammatory diseases in relation to inflammation-related carcinogenesis**, Kawanishi, S., Hiraku, Y., Pinlaor, S., Ma, N., Biol Chem. 2006 Apr;387(4):365-72.

**Nitrative and oxidative DNA damage in oral lichen planus in relation to human oral carcinogenesis**, Chaiyarit, P., Ma, N., Cancer Sci. 2005 Sep;96(9):553-9.

- **The feasibility of monitoring NF-kappaB associated cytokines: Tumor Necrosis Factor Alpha(TNF-a), IL-1alpha, IL-6, and IL-8 in whole saliva for the malignant transformation of oral lichen planus**; Rhodus, N.L., Cheng, B., et al, Mol Carcinog. 2005 Oct;44(2):77-82.

- **Cytologic and DNA-cytometric examination of oral lesions in lichen planus**, Maraki, D., Becker, J. et al, J Oral Pathol Med. 2006 Apr;35(4):227-32.

- Evaluation of proliferative potential in oral lichen planus and oral lichenoid lesions using immunohistochemical expression of p53 and Ki67, Acay, R.R., Felizzola, C.R., et al, Oral Oncol. 2006 May;42(5):475-80. Epub 2005 Dec 20;

**Valuation of exfoliative cytology as prediction factor in oral mucosa lesions**, Brunotto, M., Zarate, A.M., et al, Med Oral Patol Oral Cir Bucal. 2005 Jul 1;10 Suppl 2:E92-102.

- **A preliminary investigation of an association between dental restorations and carcinoma of the tongue**, Ma, R., Epstein, J.B., et al, Eur J Cancer B Oral Oncol. 1995 Jul;31B (4):232-4.

- **Field cancerization in oral lichen planus**, Mignogna MD, Fedele S, et al, Eur J Surg Oncol. 2006 Nov 2.

- **Two siblings with lichen planus and squamous cell carcinoma of the oesophagus**, Schwartz, M.P., Sigurdsson, V., et al, Eur J Gastroenterol Hepatol. 2006 Oct;18 (10):1111-5.

- **Oro-vaginal-vulvar lichen planus: report of two new cases**, Petruzzi, M., De, Benedittis, M., et al, Maturitas. 2005 Feb 14; 50(2):140-50.

**Unexpectedly high frequency of genital involvement in women with clinical and histological features of oral lichen planus**, Di Fede O, Belfiore, P., et al, Acta Derm Venereol. 2006;86(5):433-8.

- **Influence of treatment of erosive lichen planus of the vulva on its prognosis**, Cooper, S.M., Wojnarowska, F., Arch Dermatol. 2006 Mar;142(3):289-94.

**A clinical study of 23 cases of female genital carcinoma**, Derrick, E.K., Ridley, C.M., et al, Br J Dermatol. 2000 Dec;143(6):1217-23.

- **Note:** Etc. in a list denotes that author is aware of more references on this subject, generally available in (41 or 43).